

Copyright © 2024 by the Non-Hospital Medical and Surgical Facilities Accreditation Program and the College of Physicians and Surgeons of British Columbia.

All rights reserved. No part of this publication may be used, reproduced or transmitted, in any form or by any means electronic, mechanical, photocopying, recording or otherwise, or stored in any retrieval system or any nature, without the prior written permission of the copyright holder, application for which shall be made to:

Non-Hospital Medical and Surgical Facilities Accreditation Program College of Physicians and Surgeons of British Columbia 300–669 Howe Street Vancouver BC V6C 0B4

The Non-Hospital Medical and Surgical Facilities Accreditation Program and the College of Physicians and Surgeons of BC has used their best efforts in preparing this publication. As websites are constantly changing, some of the website addresses in this publication may have moved or no longer exist.

Introduction

Latex allergy management requires creating awareness of the potential for latex allergy and taking the appropriate steps to provide a latex-safety environment for patients. The sensitivity to latex varies from patient to patient and symptoms can range from mild (e.g. redness) to severe (e.g. anaphylaxis). Latex sensitivity symptoms can also become worse with repeated exposure. Therefore, minimizing latex exposure for latex sensitive patients is essential.

No.	Description
ALGY1.0	LATEX ALLERGY
ALGY1.1	Patients with a known or suspected latex sensitivity or allergy are identified.
ALGY1.1.1	M All patients are assessed preoperatively for latex sensitivity risk factors.
ALGY1.1.2	M Patients with a sensitivity or allergy to latex are identified with an allergy alert wristband.
ALGY1.1.3	M The medical record of any patient with a sensitivity or allergy to latex is labelled or flagged.
ALGTT.T.3	in the medical record of any patient with a sensitivity of affergy to latex is labelled of hagged.
ALGY1.1.4	M The patient's sensitivity or allergy status is clearly documented throughout the medical record.
ALGY1.2	Datients with a known or suspected latey consitivity or allergy are appropriately managed
	Patients with a known or suspected latex sensitivity or allergy are appropriately managed.
ALGY1.2.1	M Lutex productions are implemented for patients with a sensitivity of allergy to latex. 08 3 3 4 66 13 4 4



Latex Allergy

Document ID: 10826

Publication date: 2024-11-22 Version: 2.3

College of Physicians and Surgeons of British Columbia

References

American Association of Nurse Anesthetists. Latex allergy management guidelines [Internet]. Park Ridge, IL: American Association of Nurse Anesthetists; 2014. [cited 2018 May 4]. 7 p.

American Latex Allergy Association. Resources and tools: AANA latex protocol [Internet]. Slinger, WI: American Latex Allergy Association; 1998 [cited 2018 May 4].

American Latex Allergy Association. About latex allergy: cross-reactive food [Internet]. Slinger, WI: American Latex Allergy Association; [2006] [cited 2018 May 4].

Australasian Society of Clinical Immunology and Allergy. HP latex allergy: ASCIA guidelines for hospital management of latex allergic patients [Internet]. Brookvale NSW: Australasian Society of Clinical Immunology and Allergy; 2010. [cited 2018 May 4].

Operating Room Nurses Association of Canada (ORNAC). Standards guidelines and position statements for perioperative registered nursing practice. 13th ed. Halifax: ORNAC; 2017. Section 4.13: Latex sensitivity/allergy; p. 397-402.

Sussman G, Gold M. Guidelines for the management of latex allergies and safe latex use in health care facilities [Internet]. Arlington Heights, IL: American College of Allergy, Asthma & Immunology; 1996 [cited 2018 May 4].



Latex Allergy Version: 2.3

Effective date: September 21, 2018